


Figure 5.—Percent distribution of 130 brands of cigarettes and 25 brands of little cigars by nicotine content.



SOURCE: U.S. Department of Health, Education, and Welfare (97) and Federal Trade Commission (34).

TABLE 37.—*Shipment of small and large cigars destined for domestic consumption (1970, 1971, 1972)*

Year	1970	1971	1972
Small cigars			
January.....	58,328,520	85,753,780	123,477,550
February.....	63,431,580	72,092,205	179,817,839
March.....	85,881,860	46,542,800	198,165,593
April.....	101,613,500	59,059,920	125,335,740
May.....	81,093,180	93,237,473	159,334,565
June.....	82,471,120	94,560,140	180,582,243
Subtotal.....	472,819,760	451,246,318	966,713,530
July.....	62,143,140	70,332,500	127,713,320
August.....	68,220,365	127,709,310	670,936,869
September.....	79,101,045	95,027,340	422,534,705
October.....	90,752,880	109,567,900	708,116,830
November.....	64,290,600	106,666,107	551,326,888
December.....	63,806,010	123,809,553	485,587,014
Subtotal.....	428,314,040	633,112,710	2,966,215,626
Yearly total.....	901,133,800	1,084,359,028	3,932,929,156
Large cigars			
January.....	581,742,001	573,039,120	534,565,488
February.....	595,249,522	586,810,844	562,414,577
March.....	629,977,375	665,998,099	654,827,796
April.....	652,800,200	655,850,213	554,242,048
May.....	748,040,796	670,064,933	719,489,529
June.....	644,539,031	692,436,529	578,501,068
Subtotal.....	3,852,348,925	3,844,199,738	3,604,040,506
July.....	647,397,547	619,838,386	520,873,339
August.....	673,082,971	662,970,148	682,331,630
September.....	721,561,449	680,476,418	594,843,957
October.....	797,601,253	679,420,968	693,150,668
November.....	696,526,464	742,948,802	650,746,540
December.....	596,244,159	516,879,415	437,429,996
Subtotal.....	4,132,413,843	3,902,534,137	3,579,356,130
Yearly total.....	8,084,762,768	7,746,733,875	7,183,396,636

Source: U.S. Department of the Treasury (101).

TABLE 38.—*Selected compounds in mainstream smoke*

Smoke compound	Cigarette A (nonfilter)	Cigarette B (filter)	Little cigar A	Little cigar B	Small cigar C
"Tar", milligram per cigarette...	36. 1	20. 3	17. 4	31. 8	40. 6
Nicotine, milligram per cigarette...	2. 7	1. 4	. 6	1. 8	3. 1
Carbon monoxide, volume per- cent.....	4. 6	4. 5	5. 3	11. 1	7. 7
Carbon dioxide, volume percent...	9. 4	9. 6	8. 5	13. 2	12. 7
Reducing sugars, percent of tobacco weight.....	9. 3	7. 9	1. 5	2. 9	2. 7
Hydrogen cyanide, microgram per cigarette.....	536. 0	361. 0	381. 0	697. 0	1, 029. 0
Acetaldehyde, microgram per cigarette.....	770. 0	774. 0	630. 0	1, 238. 0	1, 150. 0
Acrolein, microgram per cigar- ette.....	105. 0	71. 0	41. 0	54. 0	66. 0
Total pyridines, micrograms per cigarette.....	82. 8	27. 3	58. 0	85. 3	80. 3
Phenol, microgram per cigarette...	124. 2	33. 0	35. 1	63. 4	94. 1
Benz(a)anthracene, nanogram per cigarette.....	74. 0	31. 0	34. 0	25. 0	39. 0
Benzo(a)pyrene, nanogram per cigarette.....	47. 0	20. 0	18. 0	22. 0	30. 0

Source: Hoffmann, D., Wynder, E. L. (44).

TABLE 39.—*The pH of the mainstream smoke of selected tobacco products*

[Numbers in parentheses indicate number of last puff.]

Average pH	Cigarette A (nonfilter)	Cigarette B (filter)	Little cigar A	Little cigar B	Small cigar C	Cigar D
3d puff.....	6. 19	6. 15	6. 44	6. 55	6. 53	6. 47
5th puff.....	6. 14	6. 12	6. 34	6. 46	6. 49	-----
7th puff.....	6. 09	6. 01	7. 03	6. 51	6. 56	-----
9th puff.....	6. 02	5. 83	-----	6. 98	6. 59	6. 27
13th puff....	-----	-----	-----	-----	-----	6. 39
18th puff....	-----	-----	-----	-----	-----	6. 41
23d puff....	-----	-----	-----	-----	-----	6. 81
28th puff....	-----	-----	-----	-----	-----	7. 22
33d puff....	-----	-----	-----	-----	-----	7. 53
38th puff....	-----	-----	-----	-----	-----	7. 78
Last puff....	5. 96(11)	5. 76(10)	7. 73 (8)	7. 25(10)	7. 11(11)	7. 96(43)

Source: Hoffmann, D., Wynder, E. L. (44).

Conclusions

Pipe and cigar smokers in the United States as a group experience overall mortality rates that are slightly higher than those of nonsmokers, but these rates are substantially lower than those of cigarette smokers. This appears to be due to the fact that the total exposure to smoke that a pipe or cigar smoker receives from these products is relatively low. The typical cigar smoker smokes fewer than five cigars a day and the typical pipe smoker smokes less than 20 pipefuls a day. Most pipe and cigar smokers report that they do not inhale the smoke. Those who do inhale, inhale infrequently and only slightly. As a result, the harmful effects of cigar and pipe smoking appear to be largely limited to increased death rates from cancer at those sites which are exposed to the smoke of these products. Mortality rates from cancer of the oral cavity, intrinsic and extrinsic larynx, pharynx, and esophagus are approximately equal in users of cigars, pipes, and cigarettes. Inhalation is evidently not necessary to expose these sites to tobacco smoke. Although these are serious forms of cancer, they account for only about 5 percent of the cancer mortality among men.

Coronary heart disease, lung cancer, emphysema, chronic bronchitis, cancer of the pancreas, and cancer of the urinary bladder are diseases which are clearly associated with cigarette smoking, but for cigar and pipe smokers death rates from these diseases are not greatly elevated above the rates of nonsmokers. These diseases seem to depend on moderate to deep inhalation to bring the smoke into direct contact with the issue at risk or to allow certain constituents, such as carbon monoxide, to be systematically absorbed through the lungs or to affect the temporal patterns of absorption of other constituents such as nicotine that can be absorbed either through the oral mucosa or through the lungs. Evidence from countries where smokers tend to consume more cigars and inhale them to a greater degree than in the United States indicates that rates of lung cancer become elevated to levels approaching those of cigarette smokers.

Available data on the chemical constituents of cigar, pipe, and cigarette smoke suggest that there are marked similarities in the composition of these products. Pipe and cigar smoke, however, tends to be more alkaline than cigarette smoke, and fermented tobaccos commonly used in pipes and cigars contain less reducing sugars than the rapidly dried varieties commonly used in cigarettes.

Experimental evidence suggests that little difference exists between the tumorigenic activities of tars obtained from cigar or cigarette

tobaccos. Malignant skin tumors appear somewhat more rapidly and in larger numbers in animals whose skin has been painted with cigar tars than in those animals painted with cigarette tars.

One must conclude that some risk exists from smoking cigars and pipes as they are currently used in the United States, but for most diseases this is small compared to the risk of smoking cigarettes as they are commonly used. Nevertheless, changes in patterns of usage that would bring about increased exposure either through increased individual use of cigars and pipes or increased inhalation of pipe and cigar smoke have the potential of producing risks not unlike those now incurred by cigarette smokers. Mechanical or chemical modifications of pipe tobacco and cigars that would result in a smoke more compatible with inhalation could have this effect.

Pipe and Cigar References

- (1) ABELIN, T., GSELL, O. T. Relative risk of pulmonary cancer in cigar and pipe smokers. *Cancer* 20(8) : 1288-1296, August 1967.
- (2) ALLIBONE, A., FLINT, F. J. Bronchitis, aspirin, smoking, and other factors in the aetiology of peptic ulcer. *Lancet* 2 : 179-182, July 26, 1958.
- (3) ARMITAGE, A. K., TURNER, D. M. Absorption of nicotine in cigarette and cigar smoke through the oral mucosa. *Nature* 226(5252) : 1231-1232, June 27, 1970.
- (4) ASHFORD, J. R., BROWN, S., DUFFIELD, D. P., SMITH, C. S., FAY, J. W. J. The relation between smoking habits and physique, respiratory symptoms, ventilatory function, and radiological pneumoconiosis amongst coal workers at three Scottish collieries. *British Journal of Preventive and Social Medicine* 15 : 106-117, 1961.
- (5) AUERBACH, O., HAMMOND, E. C., GARFINKEL, L. Histologic changes in the larynx in relation to smoking habits. *Cancer* 25(1) : 92-104, January 1970.
- (6) AUERBACH, O., STOUT, A. P., HAMMOND, E. C., GARFINKEL, L. Changes in bronchial epithelium in relation to sex, age, residence, smoking and pneumonia. *New England Journal of Medicine* 267(3) : 111-119, July 19, 1962.
- (7) AUERBACH, O., STOUT, A. P., HAMMOND, E. C., GARFINKEL, L. Histologic changes in esophagus in relation to smoking habits. *Archives of Environmental Health* 11(1) : 4-15, July 1965.
- (8) AUERBACH, O., STOUT, A. P., HAMMOND, E. C., GARFINKEL, L. Smoking habits and age in relation to pulmonary changes. Rupture of alveolar septums, fibrosis and thickening of walls of small arteries and arterioles. *New England Journal of Medicine* 269(20) : 1045-1054, Nov. 14, 1963.
- (9) BEST, E. W. R. A Canadian Study of Smoking and Health. Ottawa, Department of National Health and Welfare, 1966, 137 pp.
- (10) BOAKE, W. C. A study of illness in a group of Cleveland families. XVIII. Tobacco smoking and respiratory infections. *New England Journal of Medicine* 259(26) : 1245-1249, Dec. 25, 1958.
- (11) BOWER, G. Respiratory symptoms and ventilatory function in 172 adults employed in a bank. *American Review of Respiratory Diseases* 83 : 684-689, 1961.

- (12) BRADSHAW, E., SCHONLAND, M. Oesophageal and lung cancers in Natal African males in relation to certain socioeconomic factors. An analysis of 484 interviews. *British Journal of Cancer* 23(2) : 275-284, June 1969.
- (13) BRODEURS, A. C. Squamous-cell epithelioma of the lip. A study of 537 cases. *Journal of the American Medical Association* 74: 656-664, Mar. 6, 1920.
- (14) BROSS, I. D. J., TIDINGS, J. Switching from cigarettes to small cigars—is it likely to reduce the health hazards of smoking? *In press*: Proceedings of the American Association for Cancer Research, March 1973.
- (15) BRUNNEMANN, K. D., HOFFMANN, D. On the pH of tobacco smoke. Paper presented at the CORESTA/TCRC meeting, Williamsburg, Va., Oct. 23, 1972, 18 pp.
- (16) BUELL, P., DUNN, J. E., JR., BRESLOW, L. Cancer of the lung and Los Angeles-type air pollution. Prospective study. *Cancer* 20(12) : 2139-2147, December 1967.
- (17) CAMPBELL, J. M., LINDSEY, A. J. Polycyclic hydrocarbons in cigar smoke. *British Journal of Cancer* 11: 192-195, 1957.
- (18) CEDERLÖF, R., FRIBERG, L., HRUBEC, Z. Cardiovascular and respiratory symptoms in relation to tobacco smoking. A study on American twins. *Archives of Environmental Health* 18(6) : 934-940, June 1969.
- (19) COMSTOCK, G. W., BROWNLOW, W. J., STONE, R. W., SARTWELL, P. E. Cigarette smoking and changes in respiratory findings. *Archives of Environmental Health* 21(1) : 50-57, July 1970.
- (20) CRONINGER, A. B., GRAHAM, E. A., WYNDER, E. L. Experimental production of carcinoma with tobacco products. V. Carcinoma induction in mice with cigar, pipe, and all-tobacco cigarette tar. *Cancer Research* 18: 1263-1271, December 1958.
- (21) DALHAMN, T., RYLANDER, R. Ciliotoxicity of cigar and cigarette smoke. *Archives of Environmental Health* 20(2) : 252-253, February 1970.
- (22) DAVIES, R. F., DAY, T. D. A study of the comparative carcinogenicity of cigarette and cigar smoke condensate on mouse skin. *British Journal of Cancer* 23(2) : 363-368, June 1969.
- (23) DAWBER, T. R., KANNEL, W. B., REVOTSKIE, N., STOKES, J., KAGAN, A., GORDON, T. Some factors associated with the development of coronary heart disease. Six years' followup experience in the Framingham study. *American Journal of Public Health and the Nation's Health* 49(10) : 1349-1356, October 1959.
- (24) DENSEN, P. M., JONES, E. W., BASS, H. E., BREUER, J. A survey of respiratory disease among New York City postal and transit workers. I. Prevalence of symptoms. *Environmental Research* 1(3) : 262-286, November 1967.
- (25) DOLL, R., HILL, A. B. A study of the aetiology of carcinoma of the lung. *British Medical Journal* 2: 1271-1286, Dec. 13, 1952.
- (26) DOLL, R., HILL, A. B. Mortality in relation to smoking: Ten years' observations of British doctors. (Part 1). *British Medical Journal* 1(5395) : 1399-1410, May 30, 1964.
- (27) DOLL, R., HILL, A. B. Mortality in relation to smoking: Ten years' observations of British doctors. (Concluded). *British Medical Journal* 1(5396) : 1460-1467, June 6, 1964.
- (28) DOLL, R., HILL, A. B. Mortality of British doctors in relation to smoking: Observations on coronary thrombosis. *In*: Haenszel, W. (Editor). *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. Bethesda, Md., U.S. Public Health Service, National Cancer Institute Monograph No. 19, January 1966, pp. 205-215.
- (29) DOLL, R., JONES, F. A., PYGOTT, F. Effect of smoking on the production and maintenance of gastric and duodenal ulcers. *Lancet* 1: 657-662, Mar. 29, 1958.

- (30) DOYLE, J. T., DAWBER, T. R., KANNEL, W. B., KINCH, S. H., KAHN, H. A. The relationship of cigarette smoking to coronary heart disease. The second report of the combined experience of the Albany N.Y., and Framingham, Mass., studies. *Journal of the American Medical Association* 190(10) : 886-890, Dec. 7, 1964.
- (31) DUNN, J. E., JR., LINDEN, G., BRESLOW, L. Lung cancer mortality experience of men in certain occupations in California. *American Journal of Public Health and the Nation's Health* 50(10) : 1475-1487, October 1960.
- (32) EBENIUS, B. Cancer of the lip. A clinical study of 778 cases with particular regard to predisposing factors and radium therapy. *Acta Radiologica* (Supplement 48) : 1-232, 1943.
- (33) EDWARDS, F., McKEOWN, T., WHITFIELD, A. G. W. Association between smoking and disease in men over sixty. *Lancet* 1: 196-200, Jan. 24, 1959.
- (34) FEDERAL TRADE COMMISSION. Report of tar and nicotine content of the smoke of 25 varieties of small cigars. *In: Federal Trade Commission News*, 1-0703. Jul 3 1972, 2 pp.
- (35) FRANKENBURG, W. G. Chemical changes in the harvested tobacco leaf. I. Chemical and enzymic conversions during the curing process. *Advances in Enzymology* 6: 309-387, 1946.
- (36) FRANKENBURG, W. G. Chemical changes in the harvested tobacco leaf. II. Chemical and enzymic conversions during fermentation and aging. *Advances in Enzymology* 10: 325-441, 1950.
- (37) GOLDSMITH, J. R., HECHTER, H. H., PERKINS, N. M., BORHANI, N. O. Pulmonary function and respiratory findings among longshoremen. *American Review of Respiratory Diseases* 86(6) : 867-874, December 1962.
- (38) HAMMOND, E. C. Smoking in relation to the death rates of one million men and women. *In: Haenszel, W. (Editor). Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. Bethesda, Md., U.S. Public Health Service, National Cancer Institute Monograph No. 19, January 1966, pp. 127-204.
- (39) HAMMOND, E. C., GARFINKEL, L. The influence of health on smoking habits. *In: Haenszel, W. (Editor). Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. Bethesda, Md., U.S. Public Health Service, National Cancer Institute Monograph No. 19, January 1966, pp. 269-285.
- (40) HAMMOND, E. C., HORN, D. Smoking and death rates—Report on 44 months of followup of 187,783 men. I. Total mortality. *Journal of the American Medical Association* 166(10) : 1159-1172, Mar. 8, 1958.
- (41) HAMMOND, E. C., HORN, D. Smoking and death rates—Report on 44 months of followup of 187,783 men. II. Death rates by cause. *Journal of the American Medical Association* 166(11) : 1294-1308, Mar. 15, 1958.
- (42) HIRAYAMA, T. Smoking in relation to the death rates of 265,118 men and women in Japan. A report of 5 years of followup. Presented at the American Cancer Society's Fourteenth Science Writers' Seminar, Clearwater Beach, Fla., Mar. 27, 1972, 15 pp.
- (43) HOFFMANN, D., RATHKAMP, G., WYNDER, E. L. Comparison of the yields of several selected components in the smoke from different tobacco products. *Journal of the National Cancer Institute* 31(3) : 627-637, September 1963.
- (44) HOFFMANN, D., WYNDER, E. L. Smoke of cigarettes and little cigars. An analytical comparison. *Science* 178(4066) : 1197-1199, Dec. 15, 1972.
- (45) HOMBURGER, F., TREGER, A., BAKER, J. R. Mouse-skin painting with smoke condensates from cigarettes made of pipe, cigar, and cigarette tobaccos.

Journal of the National Cancer Institute 31(6) : 1445-1459, December 1963.

- (46) HOOD, B., TIBBLIN, G., WELIN, G., ORNDAHL, G., KORSAN-BENGTSSEN, K. Myocardial infarction in early age. III. Coronary risk factors and their deficient control. *Acta Medica Scandinavica* 185(4) : 241-251, April 1969.
- (47) INTER-SOCIETY COMMISSION FOR HEART DISEASE RESOURCES. Atherosclerosis Study Group and Epidemiology Study Group. Primary prevention of the atherosclerotic diseases. *Circulation* 42(6) ; A-54-A-95, December 1970.
- (48) ISAAC, P. F., RAND, M. J. Cigarette smoking and plasma levels of nicotine. *Nature* 236(5345) :308-310, April 7, 1972.
- (49) JENKINS, C. D., ROSENMAN, R. H., ZYZANSKI, S. J. Cigarette smoking. Its relationship to coronary heart disease and related risk factors in the Western Collaborative Group Study. *Circulation* 38(6) : 1140-1155, December 1968.
- (50) KAHN, H. A. The Dorn study of smoking and mortality among U.S. veterans: Report on 8½ years of observation. In: Haenszel, W. (Editor). *Epidemiological Approaches to the Study of Cancer and Other Chronic Diseases*. Bethesda, Md., U.S. Public Health Service, National Cancer Institute Monograph No. 19, January 1966, pp. 1-125.
- (51) KELLER, A. Z. Cellular types, survival, race, nativity, occupations, habits, and associated diseases in the pathogenesis of lip cancers. *American Journal of Epidemiology* 91(5) : 486-499, May 1970.
- (52) KELLER, A. Z. Cirrhosis of the liver, alcoholism, and heavy smoking associated with cancer of the mouth and pharynx. *Cancer* 20(6) : 1015-1022, June 1967.
- (53) KENSLE, C. J. The pharmacology of tobacco smoke effects of chronic exposure. In: James, G., Rosenthal, T. (Editors). *Tobacco and Health*, Springfield, C. C. Thomas, 1962, pp. 5-20.
- (54) KERSHBAUM, A. A. comparative study of cigarette, cigar, and pipe smoking effects on blood lipids, catecholamine excretion, and nicotine content of the urine. *Acta Cardiologica* 23(4) : 317-329, 1968.
- (55) KERSHBAUM, A., BELLET, S. Cigarette, cigar, and pipe smoking. Some differences in biochemical effects. *Geriatrics* 23(3) : 126-134, March 1968. 255-260, 1953.
- (56) KOULUMIES, M. Smoking and pulmonary carcinoma. *Acta Radiologica* 39 : 255-260, 1953.
- (57) KNUDTSON, K. P. The pathologic effects of smoking tobacco on the trachea and bronchial mucosa. *American Journal of Clinical Pathology* 33(4) : 310-317, April 1960.
- (58) KUHN, H. A comparative study of cigarette and cigar smoke. In: *Proceedings of the Fourth International Tobacco Scientific Congress*. The National Tobacco Board of Greece, Athens, Sept. 19-26, 1966, pp. 967-971.
- (59) LEFCOE, N. M., WONNACOTT, T. H. The prevalence of chronic respiratory disease in the male physicians of London, Ontario. *Canadian Medical Association Journal* 102 : 381-385, Feb. 28, 1970.
- (60) LEVIN, M. L., GOLDSTEIN, H., GERHARDT, P. R. Cancer and tobacco smoking. A preliminary report. *Journal of the American Medical Association* 143(4) : 336-338. May 27, 1950.
- (61) LOMBARD, H. L., SNEGIREFF, L. S. An epidemiological study of lung cancer. *Cancer* 12(2) : 406-413, March-April 1959.
- (62) MARTINEZ, I. Factors associated with cancer of the esophagus, mouth, and pharynx in Puerto Rico. *Journal of the National Cancer Institute* 42(6) : 1069-1094, June 1969.

- (63) MARTINEZ, I. Retrospective and prospective study of carcinoma of the esophagus, mouth, and pharynx in Puerto Rico. *Boletín de la Asociación Médica de Puerto Rico* 62(6) : 170-178, June 1970.
- (64) MILLS, C. A., PORTER, M. M. Tobacco smoking and automobile-driving stress in relation to deaths from cardiac and vascular causes. *American Journal of the Medical Sciences* 234 : 35-43, July 1957.
- (65) MILLS, C. A., PORTER, M. M. Tobacco smoking habits and cancer of the mouth and respiratory system. *Cancer Research* 10 : 539-542, 1950.
- (66) MILLS, C. A., PORTER, M. M. Tobacco smoking, motor exhaust fumes, and general air pollution in relation to lung cancer incidence. *Cancer Research* 17 : 981-990, 1957.
- (67) NAEYE, R. L. Structural features in Appalachian coal workers, *In*: Key, M. M., Kerr, L. E., Bundy, M. (Editors). *Pulmonary Reactions to Coal Dust. A Review of U.S. Experience*. New York, Academic Press, 1971, pp. 93-110.
- (68) OSMAN, S., BARSON, J. Hydrocarbons of cigar smoke. *Tobacco* 159 (24) : 30-32, Dec. 11, 1964.
- (69) OSMAN, S., SCHMELTZ, I., HIGMAN, H. C., STEDMAN, R. L. Volatile phenols of cigar smoke. *Tobacco* 157(9) : 30-32, Aug. 30, 1963.
- (70) PASSEY, R. D., BLACKMORE, M. Biological effects of cigar and cigarette smoke. *British Empire Cancer Campaign for Research, Annual Report* 44(Part 2) : 6, 1966.
- (71) PASSEY, R. D., BLACKMORE, M. The causation of lung cancer. Some experimental biological effects of cigar and cigarette smoke. (Abstract) *Thorax* 22(3) : 290, May 1967.
- (72) PASSEY, R. D., BLACKMORE, M., WARBRICK-SMITH, D., JONES, R. Smoking risks of different tobaccos. *British Medical Journal* 4(5781) : 198-201, Oct. 23, 1971.
- (73) PERNU, J. An epidemiological study on cancer of the digestive organs and respiratory system. A study based on 7,078 cases. *Annales Medicinae Internae Fenniae* 49(Supplement 33) : 1960, 117 pp.
- (74) RANDIG, K. Untersuchungen zur Ätiologie des Bronchialkarzinoms. (Investigations on the etiology of bronchial carcinoma.) *Öffentliche Gesundheitsdienst* 16(9) : 305-313, December 1954.
- (75) RIMINGTON, J. Chronic bronchitis, smoking and social class. A study among working people in the towns of Mid and East Cheshire. *British Journal of Diseases of the Chest* 63(4) : 193-205, October 1969.
- (76) ROE, F. J. C., CLACK, J. C., BISHOP, D., PETO, R. Comparative carcinogenicity for mouse-skin of smoke condensates prepared from cigarettes made from the same tobacco cured by two processes. *British Journal of Cancer* 24(1) : 107-121, March 1970.
- (77) SADOWSKY, D. A., GILLIAM, A. G., CORNFELD, J. The statistical association between smoking and carcinoma of the lung. *Journal of the National Cancer Institute* 13 : 1237-1258, 1953.
- (78) SANDERUD, K. Squamous metaplasia of the respiratory tract epithelium; an autopsy study of 214 cases. II. Relation to tobacco smoking, occupation and residence. *Acta Pathologica et Microbiologica Scandinavica* 43 : 47-61, 1958.
- (79) SCHIEVELBEIN, H., EBERHARDT, R. Cardiovascular actions of nicotine and smoking. *Journal of the National Cancer Institute* 48(6) : 1785-1794, June 1972.
- (80) SCHIMMLER, W., NEFF, C., SCHIMERT, G. Risikofaktoren und Herzinfarkt. Eine retrospektive Studie. (Risk factors and myocardial infarct. A retrospective study.) *Münchener Medizinische Wochenschrift* 110(27) : 1585-1594, July 5, 1968.

- (81) SCHREK, R., BAKER, L. A., BALLARD, G. P., DOLGOFF, S. Tobacco smoking as an etiologic factor in disease. I. Cancer. *Cancer Research* 16: 49-58, 1950.
- (82) SCHWARTZ, D., DENOIX, P.-F., L'enquete Francaise sur l'etiologie du cancer broncho-pulmonaire. Role du tabac. (French investigation on the etiology of bronchopulmonary cancer. Role of tobacco.) *Semaine des Hopitaux de Paris* 33(62/7): 3630-3643, Oct. 30, 1957.
- (83) SCHWARTZ, D., DENOIX, P.-F., ANGUERA, G. Recherche des localisations du cancer associees aux facteurs tabac et alcool chez l'homme. (Research on the localizations of cancer associates with tobacco and alcoholic factors in man.) *Bulletin de l'Association Francaise pour l'Etude du Cancer* 44: 336-361, 1957.
- (84) SCHWARTZ, D., FLAMANT, R., LELLOUCH, J., DENOIX, P.-F. Results of a French survey on the role of tobacco, particularly inhalation, in different cancer sites. *Journal of the National Cancer Institute* 26(5): 1085-1108, May 1961.
- (85) SHAPIRO, S., WEINBLATT, E., FRANK, C. W., SAGER, R. V. Incidence of coronary heart disease in a population insured for medical care (HIP). Myocardial infarction, angina pectoris, and possible myocardial infarction. *American Journal of Public Health and the Nation's Health* 59(6, Supplement Part 2), June 1969, 101 pp.
- (86) SPAIN, D. M., NATHAN, D. J. Smoking habits and coronary atherosclerotic heart disease. *Journal of the American Medical Association* 177(10): 683-688, Sept. 9, 1961.
- (87) STASZEWSKI, J. Palenie a rak wargi, jamy ustnej, migdalkow I krtani. (Tobacco smoking and its relation to cancer of the mouth, tonsils and larynx.) *Nowotwory* 10(2): 121-132, 1960.
- (88) STELL, P. M. Smoking and laryngeal cancer. *Lancet* 1(7751): 617-618, Mar. 18, 1972.
- (89) STOCKS, P. Cancer incidence in North Wales and Liverpool region in relation to habits and environment. *British Empire Cancer Campaign 35th Annual Report, Supplement to Part II, 1957*, 156 pp.
- (90) SVOBODA, V. An analysis of some possible epidemiological factors involved in carcinoma of the larynx. *Neoplasma* 15(6): 677-684, 1968.
- (91) TIBBLIN, G. High blood pressure in men aged 50—A population study of men born in 1913. *Acta Medica Scandinavica (Supplementum 470)*: 1-84, 1967.
- (92) TODD, G. F. (Editor). *Statistics of Smoking in the United Kingdom*. London, Tobacco Research Council, Research Paper I, Fourth Edition, 1966, 103 pp.
- (93) TODD, G. F. (Editor). *Statistics of Smoking in the United Kingdom*. London, Tobacco Research Council, Research Paper I, Fifth Edition, 1969, 124 pp.
- (94) TODD, G. F. (Editor). *Statistics of Smoking in the United Kingdom*. London, Tobacco Research Council, Research Paper I, Sixth Edition, 1972, 132 pp.
- (95) TROWELL, O. A. The relation of tobacco smoking to the incidence of chronic duodenal ulcer. *Lancet* 1: 808-809, Apr. 14, 1934.
- (96) UNITED STATES CODE OF FEDERAL REGULATIONS. Title 26 [Internal Revenue Service] Part 270—Manufacture of Cigars and Cigarettes. (As adopted at 26 F.R. 8173, effective Oct. 1, 1961). B(270)3-B(270)5, Oct. 31, 1970.
- (97) U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. Tar and Nicotine Content of Cigarettes. Washington, Federal Trade Commission, May 1972. DHEW Publication No. (HSM) 72-7510, 2 pp.

- (98) U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. National Clearinghouse for Smoking and Health. Survey conducted by National Analysts, Inc., Philadelphia, Pa., fall 1964.
- (99) U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. National Clearinghouse for Smoking and Health. Survey conducted by National Analysts, Inc., Philadelphia, Pa., and Opinion Research Corp., Princeton, N.J., spring 1966.
- (100) U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE. National Clearinghouse for Smoking and Health. Survey conducted by the Chilton Research Services, Philadelphia, Pa., January 1970.
- (101) U.S. DEPARTMENT OF THE TREASURY, BUREAU OF ALCOHOL, TOBACCO AND FIREARMS. Statistical Release: Cigarettes and Cigars, 1970, 1971, 1972.
- (102) U.S. DEPARTMENT OF THE TREASURY. Internal Revenue Cumulative Bulletin. 1969-1. Revenue Ruling 69-198, p. 359.
- (103) VAN BUCHEM, F. S. P. Serum lipids, nutrition and atherosclerotic complications in man. *Acta Medica Scandinavica* 181(4): 403-416, April 1967.
- (104) VILLIGER, U., HEYDEN-STUCKY, S. Das Infarktprofil. Unterschiede zwischen Infarktpatienten und Kontrollpersonen in des Ostschweiz. (The infarct profile. Differences between infarct patients and control persons in East Switzerland.) *Schweizerische Medizinische Wochenschrift* 96(23): 748-758, June 11, 1966.
- (105) WEIR, J. M., DUNN, J. E., Jr. Smoking and mortality: A prospective study. *Cancer* 25(1): 105-112, January 1970.
- (106) WICKEN, A. J. Environmental and Personal Factors in Lung Cancer and Bronchitis Mortality in Northern Ireland, 1960-62. Research Paper No. 9, London, Tobacco Research Council, 1966, 84 pp.
- (107) WYNDER, E. L., BROSS, I. J. A study of etiological factors in cancer of the esophagus. *Cancer* 14(2): 389-413, March-April 1961.
- (108) WYNDER, E. L., BROSS, I. J., DAY, E. A study of environmental factors in cancer of the larynx. *Cancer* 9(1): 86-110, January-February 1956.
- (109) WYNDER, E. L., BROSS, I. J., FELDMAN, R. M. A study of the etiological factors in cancer of the mouth. *Cancer* 10(6): 1300-1323, November-December 1957.
- (110) WYNDER, E. L., CORNFIELD, J. Cancer of the lung in physicians. *New England Journal of Medicine* 248(11): 441-444, Mar. 12, 1953.
- (111) WYNDER, E. L., GRAHAM, E. A. Tobacco smoking as a possible etiologic factor in bronchiogenic carcinoma; a study of 684 proved cases. *Journal of the American Medical Association* 143(4): 328-329, May 27, 1950.
- (112) WYNDER, E. L., HOFFMANN, D. Tobacco and Tobacco Smoke. Studies in Experimental Carcinogenesis. New York, Academic Press, 1967, 730 pp.
- (113) WYNDER, E. L., HULTBERG, S., JACOBSSON, F., BROSS, I. J. Environmental factors in cancer of the upper alimentary tract. A Swedish study with special reference to Plummer-Vinson (Patterson-Kelly) syndrome. *Cancer* 10(3): 470-487, May-June 1957.
- (114) WYNDER, E. L., LEMON, F. R., MANTEL, N. Epidemiology of persistent cough. *American Review of Respiratory Diseases* 91(5): 679-700, May 1965.
- (115) WYNDER, E. L., MABUCHI, K., BEATTIE, E. J., Jr. The epidemiology of lung cancer. *Journal of the American Medical Association* 213(13): 2221-2228, Sept. 28, 1970.
- (116) WYNDER, E. L., NAVARRETTE, A., AROSTEGUI, G. E., LLAMBES, J. L. Study of environmental factors in cancer of the respiratory tract in Cuba. *Journal of the National Cancer Institute* 20(4): 665-673, April 1958.
- (117) WYNDER, E. L., WRIGHT, G. A study of tobacco carcinogenesis. I. The primary fractions. *Cancer* 10(2): 255-271. March-April 1957.

CHAPTER 7

Exercise Performance

Contents

	Page
Introduction.....	241
Studies of Smokers.....	241
Studies Comparing Smokers to Nonsmokers:	
<i>Athletic Performance</i>	243
<i>Bicycle Ergometer Performance</i>	244
<i>Treadmill Performance</i>	245
<i>Performance in Other Tests of Fitness</i>	245
Discussion.....	246
Biomechanisms.....	246
Summary.....	247
References.....	248

Introduction

Although it has long been held by athletes and coaches that cigarette smoking is associated with "shortness of wind" and impaired performance, until recently there has been little scientific evidence to support this view. In the past few years, a variety of studies have appeared dealing with the effect of cigarette smoking on the response of man to exercise. The following is a review of these studies.

Age, sex, training, health, weight, and other factors are known to influence exercise performance. Because most of the investigations were carried out in healthy, young male volunteers, the groups were quite comparable with regard to age, sex, and health; however, weight, training, and other factors were often inadequately controlled. Furthermore, problems in study design and statistical analysis limit the value of several of these studies.

Many forms of exercise were performed in these experiments, including: pedaling a bicycle ergometer, running on a treadmill, running on a track, swimming, step climbing, gripping a hand dynamometer, and doing several different exercise activities as part of a battery of tests. Small to maximum amounts of work were carried out in the various studies reviewed.

Studies of Smokers

Most of the studies of habitual cigarette smokers followed a similar format with respect to smoking: (*a*) The subjects refrained from smoking for a few hours prior to testing, and (*b*) two test runs were performed, one without smoking and one in which smoking immediately preceded the exercise or was incorporated with the exercise protocol.

Several investigators (*1, 15, 28*) studied the effect of smoking on maximum grip strength. Willgoose (*28*) reported a greater mean percent recovery of grip strength after the nonsmoking trial than after the smoking trial. Kay and Karpovich (*15*) and Anderson and Brown (*1*) all followed a protocol similar to that of Willgoose except that they randomized the smoking and nonsmoking trials, and substituted

a "placebo" cigarette for the nonsmoking trial. In neither of these studies were statistically significant differences observed between the grip scores for the smoking and nonsmoking trials.

Reeves and Morehouse (24) administered a battery of tests to 15 colleges students. The tests were: A tapping test, a strength test, a jumping test, and the short form of the Harvard step test. No statistically significant differences in performance were noted under conditions of smoking or nonsmoking.

A total of 32 college students from intermediate swimming classes abstained from smoking for 15 minutes, 2 hours, and 12 hours in a study conducted by Pleasants, et al. (23). Following the abstinence, they swam distances of 100 and 200 yards. Although actual swimming times were not published, the authors reported no statistically significant differences between the mean swimming times after the different periods of abstinence for either distance.

In 1946, Juurup and Muido (13) carried out several experiments in which three young cigarette smokers exercised on a Krogh's bicycle ergometer. Smoking was found to increase the pulse rate at rest as well as during exercise. Although the effect was less consistent than on the heart rate, smoking was also associated with elevated blood pressure. Smoking had no effect on oxygen consumption. Henry and Fitzhenry (12), in 1949, using the bicycle ergometer, also found that smoking exerted no effect on oxygen consumption. In the same year, Karpovich and Hale (14) studied bicycle ergometer performance in eight young men. In all subjects, the average riding time was better in nonsmoking tests than in smoking tests; however, the results were statistically significant for only three of the eight subjects.

Kerrigan, et al. (16) more recently measured direct arterial blood pressure, heart rate, and cardiac output in 25 habitual smokers at rest and after exercise. Smoking two cigarettes produced statistically significant ($P < 0.01$) increases in cardiac index, heart rate, and arterial mean pressure compared to the immediately preceding control period. Exercise after smoking resulted in an increase in cardiac index over either the resting period or the exercise period which followed abstinence; the resultant cardiac index appeared to be approximately the sum of the exercise and smoking effects. Exercise tests preceded by smoking were also associated with significantly higher ($P < 0.01$) and more prolonged elevations of blood pressure than those not preceded by smoking.

In the study by Goldbarg, et al. (11) of nine habitual smokers performing submaximal exercise on a bicycle ergometer, cardiovascular responses were measured via pulmonary and subclavian artery catheters. At rest, after smoking, the mean cardiac index and mean heart rate increased. During successively increasing levels of exercise, the heart rate was greater and stroke index lower than values for

comparable work before smoking. The net effect of smoking was to decrease the efficiency of the heart during exercise in the upright position by causing a smaller stroke volume and a higher heart rate.

Rode and Shephard (26) investigated near maximal treadmill exercise performance in six habitual smokers. A 1-day abstinence from cigarette smoking was associated with a 13- to 79-percent decrease in the oxygen cost of breathing. Abstinence was also followed by a slowing of the heart rate and a decrease in expiratory minute volume after exercise.

The study of Krumholz, et al. (18) is different from those cited previously in that bicycle ergometer exercise performance was measured in habitual smokers both before and after 3 to 6 weeks of abstinence. Among the 10 subjects who abstained from smoking for 3 weeks, there was a statistically significant ($P < 0.05$) decrease in heart rate, oxygen debt, and ratio of oxygen debt to total increase in oxygen uptake produced by the 5 minutes of exercise.

Using a "double 9-inch progressive step test" Rode and Shephard (25) studied several hundred participants of a smoking withdrawal clinic at the time of entry and at a 1-year followup. Among those who returned for the followup and who gave up smoking, absolute aerobic power increased insignificantly; however, the relative aerobic power diminished in both sexes among those who quit smoking because of the weight gain experienced.

Studies Comparing Smokers to Nonsmokers

Athletic Performance

In 1968 Cooper, et al. (6) evaluated 419 airmen during their initial 6 weeks on active duty in the USAF. A 12-minute maximum running test was performed at least 1 hour after cigarette smoking. The mean distance covered in 12 minutes by the nonsmokers was significantly greater ($P < 0.05$) than that covered by the smokers at the beginning, the middle, and the end of training. All categories of smokers and nonsmokers improved their performance at the end of training; however, the maximum change in performance of those smoking 10 to 30 cigarettes per day was significantly ($P < 0.001$) less than that of nonsmokers.

David (7) administered a battery of tests to 88 military personnel, aged 19 to 39 years. A 1-mile run was included in the testing, and cigarette smoking was associated with a significant decrease in performance in this event.

Some 45 special forces soldiers were investigated at sea level and 13,000 feet above sea level by Fine (8). The subjects were randomly assigned to a placebo group or an acetazolamide treated group. Cigarette smoking was positively correlated to decrements in 600-yard running performance from sea level to altitude in both groups.

Pleasants (22) studied 106 students from intermediate university swimming classes. Swimming times were measured for 100- and 200-yard distances before and after training and for 800-yard distances after training. The mean swimming times of nonsmokers were less than those of smokers in six of seven listed categories, but these differences were not statistically significant.

Bicycle Ergometer Performance

Chevalier, et al. (5) investigated cardiovascular parameters in 32 young physicians after a standard 5-minute ergometer test. Oxygen debt accumulation among smokers was significantly ($P < 0.01$) greater than among nonsmokers. The heart rate at rest and 3 minutes after exercise was significantly ($P < 0.02$) faster in smokers than in nonsmokers.

Using a 5-minute ergometer test, 18 housestaff physicians, half of whom smoked, were investigated by Krumholz, et al. (17). They noted the following: Oxygen debt accumulation after exercise was significantly ($P < 0.02$) greater in smokers than non-smokers, the ratio of the oxygen debt to total increased oxygen uptake during exercise was significantly ($P < 0.001$) greater in smokers than in nonsmokers, and the diffusing capacity at rest and with exercise was significantly ($P < 0.05$) decreased in smokers compared to nonsmokers.

Kerrigan, et al. (16) studied cardiovascular parameters in smokers and nonsmokers at rest, during, and after a 5-minute bicycle ergometer ride. Cardiac index and blood pressure values obtained during exercise performed immediately after smoking were greater than those found in nonsmokers performing the same exercise. Similarly, heart rate and blood pressure remained elevated for longer periods in those who exercised immediately after smoking than in nonsmokers performing the same task.

Aerobic capacity scores were examined in 60 university student volunteers by Peterson and Kelley (20). Subjects worked at submaximal levels on a bicycle ergometer before, during, and after a training program. At all of these intervals, nonsmokers had significantly ($P < 0.05$) higher mean aerobic capacity scores than smokers. Both groups increased their aerobic capacity during training but nonsmokers consistently performed better throughout training.

Treadmill Performance

In 1960 Blackburn, et al. (4) carried out several measurements of cardiovascular function after different amounts of treadmill exercise were performed by 233 professional men, 159 university students, and 414 railroad workers. The differences between the smokers and nonsmokers were of small magnitude. Basal oxygen consumption was slightly higher in smokers than in nonsmokers. Also, resting pulse rates were higher in smokers of most groups.

Cooper, et al. (6) studied 47 out of 419 airmen with treadmill testing. Cardiopulmonary indices measured on the treadmill, including maximum indices, were comparable in smokers and nonsmokers except for a significant ($P < 0.01$) reduction in the maximum minute volume among the smokers.

A total of 277 prospective Canadian firemen performed the Balke-Ware test of work capacity in treadmill studies carried out by Glassford and Howell (10). The mean performance scores of nonsmokers were significantly ($P < 0.01$) greater than those of smokers.

The effect of vitamin C supplementation on treadmill exercise performance was investigated in 40 male volunteers by Bailey, et al. (3). Significant differences in oxygen utilization and ventilatory function between smokers and nonsmokers were noted in only two of the 24 separate analyses of variance performed.

Maximal oxygen intake during treadmill exercise was examined by McDonough, et al. (19) in 86 healthy, middle-aged male volunteers. Cigarette smoking was one of six variables which together provided a multiple correlation coefficient of 0.73.

Performance in Other Tests of Fitness

When physical fitness tests were administered to 88 military personnel by David (7), cigarette smoking was found to be associated with a significant ($P < 0.001$) decrease in performance in the dodge and jump test, and a significant ($P < 0.02$) decrease in performance in the crawling test.

Using a step test, a breath holding test, and an ergometer test, Franks (9) examined 58 middle-aged men. Nonsmokers were able to hold their breath longer and had greater vital capacity residual after the step test than the smokers.

In 1971, Wysokinski (29) studied 200 young Polish soldiers using Letunov's test which included 20 knee-bending exercises, a fast run for 20 seconds, and a run for 3 minutes. Cigarette smoking was associated with a significant ($P < 0.01$) reduction in the vital capacity and a

marked rise in the pulse rate at rest and after exercise. Intense exercise also caused a greater rise in the systolic blood pressure in smokers than in nonsmokers.

Discussion

Most of the studies in habitual cigarette smokers compared exercise performance in "smoking" and "nonsmoking" runs after only a few hours of abstinence. In some studies, smoking adversely affected performance (11, 13, 14, 16, 18, 26, 28), while in others it did not (1, 12, 15, 23, 24). Some of these apparently discrepant results are due to differences in methodology and in amounts and types of work performed. In all of the more recent studies of habitual smokers in which moderate to near maximal amounts of work were performed and sophisticated measurements of oxygen transport and cardiopulmonary function were made, impairment of function during smoking trials was found (11, 16, 18, 26).

The data of Krumholz, et al. (18) also raise the question of whether residual effects of cigarette smoking influence "nonsmoking" trials performed after a few hours of abstinence; they found statistically significant decreases in heart rate and oxygen debt produced by exercise after 3 weeks of cessation.

The work of Rode and Shephard (25) suggests that physical fitness improves with cessation, but this improvement may be negated if the subject gains a substantial amount of weight after giving up smoking.

Several investigators compared exercise performance or postexercise cardiopulmonary function of smokers to nonsmokers. Although only minor differences between smokers and nonsmokers were found in a few of these studies (3, 4, 22), in most of them (5, 6, 7, 8, 10, 16, 17, 20, 29) the performance or function of the nonsmokers was better than that of the smokers. Both nonsmokers and smokers improved their performance with training, but nonsmokers maintained their advantage throughout training (6, 20).

Biomechanisms

The cited studies indicate that cigarette smoking exerts its adverse effect on exercise performance through several mechanisms. Cigarette smoking appears to impair cardiac performance during exercise by increasing the heart rate and exerting a variable effect on cardiac

output (5, 11, 13, 16, 18, 26, 29). Cigarette smoking is associated with an increased oxygen debt after exercise (5, 18). Also, one study indicated that the oxygen cost of hyperventilation was greater among smokers than among nonsmokers (26).

Some of these adverse effects of smoking on oxidative metabolism are mediated by the elevated carboxyhemoglobin levels found in smokers. CO exerts these effects through one or more of the following mechanisms: (a) Reduction of the amount of hemoglobin available for oxygen transport, (b) shift of the oxygen-hemoglobin dissociation curve to the left with consequent interference in oxygen release at the tissue level, (c) induction of arterial hypoxemia, and (d) possible interference with the homeostatic mechanism by which 2,3,DPG controls the affinity of hemoglobin for oxygen (27). Because carboxyhemoglobin has a half life in the body of at least 3 to 4 hours, its influence may still be measurable several hours after abstinence from smoking (27).

A recent investigation of maximal muscular exercise during CO intoxication in five male volunteers demonstrated reduced maximal O₂ consumption in spite of a much higher heart rate and a relative hyperventilation (21).

Astrand and Rodahl (2) commented recently on the adverse effect of cigarette smoking on oxygen transport: "All other factors being equal, a reduction in the oxygen-transporting capacity is associated with a corresponding reduction in physical performance capacity during heavy or maximal work * * *. Because a regular physical training program only increases the maximal oxygen uptake by some 10 to 20 percent, a 5- to 10-percent reduction in maximal aerobic power due to smoking may play a significant role in many types of athletic events and in very heavy work."

Other studies cited in this review document the adverse effect of smoking on pulmonary diffusing capacity (18) and on pulmonary function with exercise (6, 29).

Summary

Clinical studies in healthy, young men have shown that cigarette smoking impairs exercise performance, especially for many types of athletic events and activities involving maximal work capacity. Some of these effects are mediated by reduced oxygen transport and reduced cardiac and pulmonary function.

Exercise Performance References

- (1) ANDERSON, J. M., BROWN, C. W. A study of the effects of smoking upon grip strength and recuperation from local muscular fatigue. *Research Quarterly* 22(1) : 102-108, March 1951.
- (2) ASTRAND, P.-O., RODAHL, K. Factors affecting performance. In: *Textbook of Work Physiology*. New York, McGraw-Hill Book Co., 1970, 669 pp.
- (3) BAILEY, D. A., CARRON, A. V., TEECE, R. G., WEHNER H. J. Vitamin C supplementation related to physiological response to exercise in smoking and nonsmoking subjects. *American Journal of Clinical Nutrition* 23(7) : 905-912, July 1970.
- (4) BLACKBURN, H., BROZEK, J., TAYLOR, H. L. Common circulatory measurements in smokers and nonsmokers. *Circulation* 22 : 1112-1124, December 1960.
- (5) CHEVALIER, R. B., BOWERS, J. A., BONDURANT, S., ROSS, J. C. Circulatory and ventilatory effects of exercise in smokers and nonsmokers. *Journal of Applied Physiology* 18(2) : 357-360, March 1963.
- (6) COOPER, K. H., GEY, G. O., BOTTENBERG, R. A. Effects of cigarette smoking on endurance performance. *Journal of the American Medical Association* 203(3) : 189-192, Jan. 15, 1968.
- (7) DAVID, K. H. Age, cigarette smoking, and tests of physical fitness. *Journal of Applied Psychology* 52(4) : 296-298, August 1968.
- (8) FINE, B. J. Personality traits as related to symptomatology and running performance at altitude under normal and drug (acetazoleamide) conditions. *Perceptual and Motor Skills* 27 : 975-990, 1968.
- (9) FRANKS, B. D. Smoking and selected cardiovascular-respiratory measures. *Research Quarterly* 41(2) : 140-144, May 1970.
- (10) GLASSFORD, R. G., HOWELL, M. L. Smoking and physical fitness: A preliminary report. *Canadian Family Physician* 15(10) : 60-62, October 1969.
- (11) GOLDBARG, A. N., KRONE, R. J., RESNEKOV, L. Effects of cigarette smoking on hemodynamics at rest and during exercise. I. Normal subjects. *Chest* 60(6) : 531-536, December 1971.
- (12) HENRY, F. M., FITZHENRY, J. R. Oxygen metabolism of moderate exercise, with some observations on the effects of tobacco smoking. *Journal of Applied Physiology* 2 : 464-468, February 1950.
- (13) JUURUP, A., MUIDO, L. On acute effects of cigarette smoking on oxygen consumption, pulse rate, breathing rate and blood pressure in working organisms. *Acta Physiologica Scandinavica* 11 : 48-60, 1946.
- (14) KARPOVICH, P. V., HALE, C. J. Tobacco smoking and physical performance. *Journal of Applied Physiology* 3 : 616-621, April 1951.
- (15) KAY, H. W., KARPOVICH, P. V. Effect of smoking upon recuperation from local muscular fatigue. *Research Quarterly* 20 : 250-256, 1949.
- (16) KERRIGAN, R., JAIN, A. C., DOYLE, J. T. The circulatory response to cigarette smoking at rest and after exercise. *American Journal of the Medical Sciences* 255 : 113-119, February 1968.
- (17) KRUMHOLZ, R. A., CHEVALIER, R. B., ROSS, J. C. Cardiopulmonary function in young smokers. A comparison of pulmonary function measurements and some cardiopulmonary responses to exercise between a group of young smokers and a comparable group of nonsmokers. *Annals of Internal Medicine* 60(4) : 603-610, April 1964.
- (18) KRUMHOLZ, R. A., CHEVALIER, R. B., ROSS, J. C. Changes in cardiopulmonary functions related to abstinence from smoking. Studies in young cigarette smokers at rest and exercise at 3 and 6 weeks of abstinence. *Annals of Internal Medicine* 62(2) : 197-207, February 1965.

- (19) McDONOUGH, J. R., KUSUMI, F., BRUCE, R. A. Variations in maximal oxygen intake with physical activity in middle-aged men. *Circulation* 41(5) : 743-751, May 1970.
- (20) PETERSON, F. J., KELLEY, D. L. The effect of cigarette smoking upon the acquisition of physical fitness during training as measured by aerobic capacity. *Journal of the American College Health Association* 17(3) : 250-254, February 1969.
- (21) PIRNAY, F., DUJARDIN, J., DEBOANNE, R., PETIT, J. M. Muscular exercise during intoxication by carbon monoxide. *Journal of Applied Physiology* 31(4) : 573-575, October 1971.
- (22) PLEASANTS, F., JR. Pretraining and post-training swimming endurance of smokers and nonsmokers. *Research Quarterly* 40(4) : 779-782, 1969.
- (23) PLEASANTS, F., JR., GRUGAN, J., RATLIFF, J. W., JR. Effects of short periods of abstinence from cigarette smoking on swimming endurance of chronic smokers. *Research Quarterly* 38(3) : 474-479, 1966.
- (24) REEVES, W. E., MOREHOUSE, L. E. The acute effect of smoking upon the physical performance of habitual smokers. *Research Quarterly* 21 : 245-248, 1950.
- (25) RODE, A., SHEPHARD, R. J. Smoking withdrawal and changes of cardiorespiratory fitness. *American Review of Respiratory Disease* 104(6) : 933-935, December 1971.
- (26) RODE, A., SHEPHARD, R. J. The influence of cigarette smoking upon the oxygen cost of breathing in near-maximal exercise. *Medicine and Science in Sports* 3(2) : 51-55, summer 1971.
- (27) U.S. PUBLIC HEALTH SERVICE. The Health Consequences of Smoking. A Report of the Surgeon General : 1972. U.S. Department of Health, Education, and Welfare. Washington, DHEW Publication No. (HSM) 72-7516, 1972, 158 pp.
- (28) WILLGOOSE, C. E. Tobacco smoking, strength, and muscular endurance. *Research Quarterly* 18 : 219-225, 1947.
- (29) WYSOKINSKI, Z. Effects of tobacco smoking on certain parameters reflecting the condition of the circulatory system at rest and during exercise. *Polish Medical Science and History* 14(2) : 73-76, April 1971.